

Eigenvalue spectrum analysis for temporal signals of Kerr optical frequency combs based on nonlinear Fourier transform

Jia Wang(王佳), Ai-Guo Sheng(盛爱国), Xin Huang(黄鑫), Rong-Yu Li(李荣玉), Guang-Qiang He(何广强) Citation:Chin. Phys. B . 2020, 29(3): 034207 . doi: 10.1088/1674-1056/ab683a Journal homepage: <u>http://cpb.iphy.ac.cn</u>; <u>http://iopscience.iop.org/cpb</u>

What follows is a list of articles you may be interested in

Scanning the energy dissipation process of energetic materials based on excited state relaxation and vibration-vibration coupling

Wen-Yan Wang(王文岩), Ning Sui(隋宁), Li-Quan Zhang(张里荃), Ying-Hui Wang(王英惠), Lin Wang(王琳), Quan Wang(王权), Jiao Wang(王娇), Zhi-Hui Kang(康智慧), Yan-Qiang Yang(杨延强), Qiang Zhou(周强), Han-Zhuang Zhang(张 汉壮)

Chin. Phys. B . 2018, 27(10): 104205 . doi: 10.1088/1674-1056/27/10/104205

Nonlinear spectral cleaning effect in cross-polarized wave generation

Linpeng Yu(於林鹏), Yi Xu(许毅), Fenxiang Wu(吴分翔), Xiaojun Yang(杨晓骏), Zongxin Zhang(张宗昕), Yuanfeng Wu(吴圆峰), Yuxin Leng(冷雨欣), Zhizhan Xu(徐至展) Chin. Phys. B . 2018, 27(5): 054214 . doi: 10.1088/1674-1056/27/5/054214

Enhanced second harmonic generation in a two-dimensional optical micro-cavity

Jian-Jun Zhang(张建军), Hui-Fang Wang(王慧芳), Jun-Hua Hou(候俊华) Chin. Phys. B . 2018, 27(3): 034207 . **doi:** 10.1088/1674-1056/27/3/034207

Modulation instabilities in randomly birefringent two-mode optical fibers

Jin-Hua Li(李金花), Hai-Dong Ren(任海东), Shi-Xin Pei(裴世鑫), Zhao-Lou Cao(曹兆楼), Feng-Lin Xian(咸冯林) Chin. Phys. B . 2016, 25(12): 124208 . doi: 10.1088/1674-1056/25/12/124208

<u>Ultra-broadband modulation instability gain characteristics in As₂S₃ and As₂Se₃ chalcogenide glass photonic crystal fiber</u>

Ju- con

He-Lin Wang(王河林), Bin Wu(吴彬), Xiao-Long Wang(王肖隆) Chin. Phys. B . 2016, 25(6): 064207 . **doi:** 10.1088/1674-1056/25/6/064207



Volume 29 March 2020 Number 3

SPECIAL TOPIC

- Topological semimetals
- Advanced calculation & characterization of energy storage materials & devices at multiple scale

A series Journal of the Chinese Physical Society Distributed by IOP Publishing iopscience.org/cpb cpb.iphy.ac.cn



Featured Article

Tailoring electronic properties of two-dimensional antimonene with isoelectronic counterparts Ye Zhang, Huai-Hong Guo, Bao-Juan Dong, Zhen Zhu, Teng Yang, Ji-Zhang Wang, Zhi-Dong Zhang Chin, Phys. B, 2020, 29(3): 037305

Chinese Physics B(中国物理 B)

Published monthly in hard copy by the Chinese Physical Society and online by IOP Publishing, Temple Circus, Temple Way, Bristol BS1 6HG, UK

Institutional subscription information: 2020 volume

For all countries, except the United States, Canada and Central and South America, the subscription rate per annual volume is UK \pounds 1326 (electronic only) or UK \pounds 1446 (print + electronic).

Delivery is by air-speeded mail from the United Kingdom.

Orders to:

Journals Subscription Fulfilment, IOP Publishing, Temple Circus, Temple Way, Bristol BS1 6HG, UK

For the United States, Canada and Central and South America, the subscription rate per annual volume is US\$2640 (electronic only) or US\$2870 (print + electronic). Delivery is by transatlantic airfreight and onward mailing.

Orders to: IOP Publishing, P. O. Box 320, Congers, NY 10920-0320, USA

© 2020 Chinese Physical Society and IOP Publishing Ltd

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the copyright owner.

Supported by the China Association for Science and Technology and Chinese Academy of Sciences

Editorial Office: Institute of Physics, Chinese Academy of Sciences, P. O. Box 603, Beijing 100190, China Tel: (86-10) 82649026 or 82649519, Fax: (86-10) 82649027, E-mail: cpb@aphy.iphy.ac.cn

主管单位:中国科学院
主办单位:中国物理学会和中国科学院物理研究所
主编:欧阳钟灿
出版:中国物理学会
印刷装订:北京科信印刷有限公司
编辑: Chinese Physics B 编辑部
国内发行: Chinese Physics B 出版发行部
国外发行: IOP Publishing Ltd
发行范围:公开发行

国际统一刊号: ISSN 1674-1056 国内统一刊号: CN 11-5639/O4 编辑部地址: 北京 中关村 中国科学院物理研究所内 通讯地址: 100190北京603信箱 电 话: (010) 82649026, 82649519 传 真: (010) 82649027 "Chinese Physics B"网址: http://cpb.iphy.ac.cn http://iopscience.iop.org/journal/1674-1056

Published by the Chinese Physical Society

Advisory Board

Prof. Academician Chen Jia-Er(陈佳洱)

Prof. Academician Feng Duan(冯端)

Prof. Academician T. D. Lee(李政道)

Prof. Academician Samuel C. C. Ting(丁肇中)

Prof. Academician C. N. Yang(杨振宁)

Prof. Academician Yang Fu-Jia(杨福家)

Prof. Academician Zhou Guang-Zhao

(Chou Kuang-Chao)(周光召)

Prof. Academician Wang Nai-Yan(王乃彦)

Editor-in-Chief

Prof. Academician Ouyang Zhong-Can(欧阳钟灿)

School of Physics, Peking University, Beijing 100871, China Department of Physics, Nanjing University, Nanjing 210093, China Department of Physics, Columbia University, New York, NY 10027, USA LEP3, CERN, CH-1211, Geneva 23, Switzerland Institute for Theoretical Physics, State University of New York, USA Department of Nuclear Physics, Fudan University, Shanghai 200433, China China Association for Science and Technology, Beijing 100863, China

China Institute of Atomic Energy, Beijing 102413, China

Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

Associate Editors

Prof. Academician Zhao Zhong-Xian(赵忠贤)
Prof. Academician Yang Guo-Zhen(杨国桢)
Prof. Academician Zhang Jie(张杰)
Prof. Academician Xing Ding-Yu(邢定钰)
Prof. Academician Shen Bao-Gen(沈保根)
Prof. Academician Gong Qi-Huang(龚旗煌)
Prof. Academician Xue Qi-Kun(薛其坤)
Prof. Sheng Ping(沈平)

Editorial Board

Prof. David Andelman Prof. Academician Chen Xian-Hui(陈仙辉)

Prof. Cheng Jian-Chun(程建春) Prof. Chia-Ling Chien

Prof. Dai Xi(戴希) Prof. Ding Jun(丁军)

Prof. Masao Doi

Prof. Fang Zhong(方忠) Prof. Feng Shi-Ping(冯世平) Prof. Academician Gao Hong-Jun(高鸿钧) Prof. Gu Chang-Zhi(顾长志) Prof. Gu Min(顾敏)

Prof. Academician Guo Guang-Can(郭光灿)

Prof. Academician He Xian-Tu(贺贤土)

Prof. Werner A. Hofer

Prof. Hong Ming-Hui(洪明辉)

Prof. Hu Gang(胡岗) Prof. Jiang Hong-Wen(姜弘文)

Prof. Jiang Ying(江颖) Prof. Jin Xiao-Feng(金晓峰) Prof. Robert J. Joynt Prof. Jaewan Kim

Prof. Li Ru-Xin(李儒新)

Prof. Li Xiao-Guang(李晓光)

Assits. Prof. Liu Chao-Xing(刘朝星) Prof. Liu Xiang-Yang(刘向阳) Prof. Liu Ying(刘荧)

Prof. Long Gui-Lu(龙桂鲁) Prof. Lv Li(吕力) Prof. Ma Xu-Cun(马旭村) Prof. Antonio H. Castro Neto

Prof. Nie Yu-Xin(聂玉昕) Prof. Niu Qian(牛谦) Prof. Academician Ouyang Qi(欧阳颀) Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Chinese Academy of Sciences, Beijing 100864, China Department of Physics, Nanjing University, Nanjing 210093, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China School of Physics, Peking University, Beijing 100871, China Department of Physics, Tsinghua University, Beijing 100084, China The Hong Kong University of Science & Technology, Kowloon, Hong Kong, China

School of Physics and Astronomy Tel Aviv University, Tel Aviv 69978, Israel Department of Physics, University of Science and Technology of China, Hefei 230026. China School of Physics, Nanjing University, Nanjing 210093, China Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218, USA Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Materials Science & Engineering, National University of Singapore, Singapore 117576, Singapore Toyota Physical and Chemical Research Institute, Yokomichi, Nagakute, Aichi 480-1192, Japan Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Physics, Beijing Normal University, Beijing 100875, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China School of Optical-Electrical and Computer Engineering, University of Shanghai for Science and Technology, Shanghai 200093, China School of Physical Sciences, University of Science and Technology of China, Hefei 230026, China Institute of Applied Physics and Computational Mathematics, Beijing 100088, China Stephenson Institute for Renewable Energy, The University of Liverpool, Liverpool L69 3BX, UK Department of Electrical and Computer Engineering, National University of Singapore, Singapore 117576, Singapore Department of Physics, Beijing Normal University, Beijing 100875, China Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA School of Physics, Peking University, Beijing 100871, China Department of Physics, Fudan University, Shanghai 200433, China Physics Department, University of Wisconsin-Madison, Madison, USA Korea Institute for Advanced Study, School of Computational Sciences, Hoegiro 85, Seoul 02455, Korea Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, Shanghai 201800, China Department of Physics, University of Science and Technology of China, Hefei 230026, China Department of Physics, Pennsylvania State University, PA 16802-6300, USA Department of Physics, Xiamen University, Xiamen 361005, China Department of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China Department of Physics, Tsinghua University, Beijing 100084, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Physics, Tsinghua University, Beijing 100084, China Physics Department, Faculty of Science, National University of Singapore, Singapore 117546, Singapore Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Physics, University of Texas, Austin, TX 78712, USA School of Physics, Peking University, Beijing 100871, China

Prof. Academician Pan Jian-Wei(潘建伟) Prof. Amalia Patane Prof. Qian Lie-Jia(钱列加) Prof. J. Y. Rhee Prof. Shen Jian(沈健) Prof. Shen Yuan-Rang(沈元壤) Prof. Shen Zhi-Xun(沈志勋) Prof. Academician Sun Chang-Pu(孙昌璞) Prof. Sun Xiao-Wei(孙小卫) Prof. Sun Xiu-Dong(孙秀冬) Prof. Michiyoshi Tanaka Prof. Tong Li-Min(童利民) Prof. Tong Penger(童彭尔) Prof. Wang Bo-Gen(王伯根) Prof. Wang Kai-You(王开友) Prof. Wang Wei(王炜) Prof. Wang Ya-Yu(王亚愚) Prof. Wang Yu-Peng(王玉鹏) Prof. Wang Zhao-Zhong(王肇中) Prof. Academician Wang Wei-Hua(汪卫华) Prof. Wei Su-Huai(魏苏淮) Prof. Wen Hai-Hu(闻海虎) Prof. Wu Nan-Jian(吴南健) Prof. Academician Xia Jian-Bai(夏建白) Prof. Academician Xiang Tao(向涛) Prof. Academician Xie Si-Shen(解思深) Prof. Academician Xie Xin-Cheng(谢心澄) Prof. Academician Xu Zhi-Zhan(徐至展) Assist. Prof. Xu Cen-Ke(许岑珂) Prof. Academician Ye Chao-Hui(叶朝辉) Prof. Ye Jun(叶军) Prof. Yu Ming-Yang(郁明阳) Prof. Academician Zhan Wen-Long(詹文龙) Prof. Zhang Fu-Chun(张富春) Prof. Zhang Xiang(张翔) Prof. Zhang Yong(张勇) Prof. Zhang Zhen-Yu(张振宇) Prof. Zeng Hao(曾浩) Prof. Zheng Bo(郑波) Prof. Zhou Xing-Jiang(周兴江) Prof. Academician Zhu Bang-Fen(朱邦芬)

Editorial Staff

School of Physics and Astronomy, The University of Nottingham, NG7 2RD, UK Department of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China Department of Physics, Sungkyunkwan University, Suwon, Korea Department of Physics, Fudan University, Shanghai 200433, China Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA Stanford University, Stanford, CA 94305-4045, USA Beijing Computational Science Research Center, China Academy of Engineering Physics, Beijing 100094, China Department of Electrical and Electronic Engineering, Southern University of Science and Technology, Shenzhen 518055, China Department of Physics, Harbin Institute of Technology, Harbin 150001, China Research Institute for Scientic Measurements, Tohoku University, Katahira 2-1-1, Aoba-ku 980, Sendai, Japan Department of Optical Engineering, Zhejiang University, Hangzhou 310027, China Department of Physics, The Hong Kong University of Science and Technology, Kowloon, Hong Kong, China School of Physics, Nanjing University, Nanjing 210093, China Institute of Semiconductors, Chinese Academy of Sciences, Beijing 100083, China School of Physics, Nanjing University, Nanjing 210093, China Department of Physics, Tsinghua University, Beijing 100084, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Laboratory for Photonics and Nanostructures (LPN) CNRS-UPR20, Route de Nozay, 91460 Marcoussis, France Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Beijing Computational Science Research Center, China Academy of Engineering Physics, Beijing 100094, China Department of Physics, Nanjing University, Nanjing 210093, China Institute of Semiconductors, Chinese Academy of Sciences, Beijing 100083, China Institute of Semiconductors, Chinese Academy of Sciences, Beijing 100083, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Physics, Peking University, Beijing 100871, China Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, Shanghai 201800, China Department of Physics, University of California, Santa Barbara, CA 93106, USA Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China Department of Physics, University of Colorado, Boulder, Colorado 80309-0440, USA Theoretical Physics I, Ruhr University, D-44780 Bochum, Germany Chinese Academy of Sciences, Beijing 100864, China Kavli Institute for Theoretical Sciences, University of Chinese Academy of Sciences, Beijing 100190, China NSF Nanoscale Science and Engineering Center (NSEC), University of California, Berkeley, CA 94720, USA Electrical and Computer Engineering Department, The University of North Carolina at Charlotte, Charlotte, USA International Center for Quantum Design of Functional Materials, University of Science and Technology of China, Hefei 230026, China Department of Physics, University at Buffalo, SUNY, Buffalo, NY 14260, USA Physics Department, Zhejiang University, Hangzhou 310027, China Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China Department of Physics, Tsinghua University, Beijing 100084, China

Department of Modern Physics, University of Science and Technology of

China, Hefei 230026, China

Eigenvalue spectrum analysis for temporal signals of Kerr optical frequency combs based on nonlinear Fourier transform*

Jia Wang(王佳)¹, Ai-Guo Sheng(盛爱国)¹, Xin Huang(黄鑫)¹, Rong-Yu Li(李荣玉)¹, and Guang-Qiang He(何广强)^{1,2,3,†}

¹ State Key Laboratory of Advanced Optical Communication Systems and Networks, Department of Electronic Engineering, Shanghai Jiao Tong University, Shanghai 200240, China ² State Key Laboratory of Precision Spectroscopy, East China Normal University, Shanghai 200062, China ³National Laboratory of Solid State Microstructures, Nanjing University, Nanjing 210093, China

(Received 24 September 2019; revised manuscript received 20 November 2019; accepted manuscript online 7 January 2020)

Based on the nonlinear Schrödinger equation (NLSE) with damping, detuning, and driving terms describing the evolution of signals in a Kerr microresonator, we apply periodic nonlinear Fourier transform (NFT) to the study of signals during the generation of the Kerr optical frequency combs (OFCs). We find that the signals in different states, including the Turing pattern, the chaos, the single soliton state, and the multi-solitons state, can be distinguished according to different distributions of the eigenvalue spectrum. Specially, the eigenvalue spectrum of the single soliton pulse is composed of a pair of conjugate symmetric discrete eigenvalues and the quasi-continuous eigenvalue spectrum with eve-like structure. Moreover, we have successfully demonstrated that the number of discrete eigenvalue pairs in the eigenvalue spectrum corresponds to the number of solitons formed in a round-trip time inside the Kerr microresonator. This work shows that some characteristics of the time-domain signal can be well reflected in the nonlinear domain.

Keywords: nonlinear Fourier transform, Kerr optical frequency combs, nonlinear signal processing

PACS: 42.65.Sf, 42.65.Tg

1. Introduction

The nonlinear Fourier transform (NFT), also known as the inverse scattering transform (IST), is a powerful mathematical tool to solve the problems of wave propagation in some certain nonlinear media, especially in the field of soliton theory.^[1] In contrast with the linear Fourier transform (FT), where the signal is decomposed into the superposition of sine waves so as to obtain the frequency components in the linear frequency domain, the nonlinear Fourier transform can decompose the signal into soliton components and non-soliton components (also known as radiation components) to obtain the corresponding nonlinear frequency components.

In recent years, NFT has been widely used in the field of optical fiber communication. The nonlinear Schrödinger equation (NLSE) is the mathematical model to describe the signal propagation in the optical fiber channel.^[2-4] Without considering the channel noise, NLSE can be approximated as an integrable partial differential equation. In this case, the complex evolution problems of time-domain signal and linear spectrum can be converted into a simple nonlinear spectrum evolution problem by using NFT.^[5-7] Based on this characteristic, many researchers have carried out a series of theoretical studies and experiments to reduce the unfavorable restriction brought by fiber nonlinearity to optical fiber communication.^[8–12] The realization of high-speed and reliable fiber communication sys-



tem has proved the superiority of NFT in solving the NLSE problem.^[13-15] In 2018, Ryczkowski et al.^[16] used NFT to analyze the transient dissipative soliton dynamics in a modelocked laser, clearly observed the presence of local soliton content, and promoted the application of NFT in describing ultra-short pulses.

In this paper, we use NFT to study the temporal signals of Kerr optical frequency combs (OFCs) based on Kerr microresonator. The Kerr OFCs are presented as a series of precisely equidistant spectral lines with extremely narrow linewidth in the linear frequency domain, and the corresponding time-domain signals of OFCs in stable soliton state are presented as periodic ultra-short laser pulse sequences.^[17,18] We can clearly distinguish the unstable state and different stable states of OFCs, including the chaos, the Turing patterns, and various soliton states, according to the eigenvalue spectrum obtained by NFT for various temporal signals in the cavity. We also find that the number of discrete eigenvalue pairs is consistent with the number of solitons formed in the intracavity round-trip time.

A me theory 2.1. The model for Kerr OFCs generation The generation of Kerr OFC-The generation of Kerr OFCs is described by the spatiotemporal Lugiato-Lefever equation (LLE) whose normal-

*Project supported by the National Natural Science Foundation of China (Grant Nos. 61475099 and 61922040), Program of State Key Laboratory of Quantum Optics and Quantum Optics Devices, China (Grant No. KF201701), and the Key R&D Program of Guangdong Province, China (Grant No. 2018B030325002).

[†]Corresponding author. E-mail: gqhe@sjtu.edu.cn

^{© 2020} Chinese Physical Society and IOP Publishing Ltd

ized form follows^[19,20]

$$\frac{\partial \Psi}{\partial \tau} = -(1+j\zeta_0)\Psi + j|\Psi|^2\Psi - j\frac{\beta}{2}\frac{\partial^2\Psi}{\partial\phi^2} + f.$$
 (1)

On the left-hand side, $\Psi(\tau, \phi)$ is the complex envelope of the intracavity field, and τ denotes the normalized time. On the right-hand side, $\zeta_0 = 2(\omega_0 - \omega_p)/\kappa$ denotes the normalized detuning of the pump laser ω_p from the resonance frequency ω_0 , κ stands for a coupled resonance width (also known as the total linewidth), ϕ is the co-rotating angular coordinate inside the resonator, and $\beta = -2D_2/\kappa$ is the overall dispersion parameter with D_2 being the second-order dispersion, that is, the anomalous group velocity dispersion (GVD) regime follows $\beta < 0$, while the normal GVD regime is defined by $\beta > 0$. *f* represents the dimensionless external pump field intensity, for completeness, *f* can be calculated by

$$f = \sqrt{\frac{8g\eta P_{\rm in}}{\kappa^2 \hbar \omega_0}}.$$
 (2)

Here, $P_{\rm in}$ is the optical power (in W) of the laser pump at the input of the resonator, $g = \frac{\hbar \omega_0^2 c n_2}{n_0^2 V_{\rm eff}}$ is the nonlinear coupling coefficient, \hbar is the reduced Planck constant, c is the speed of light in vacuum, n_0 and n_2 are the linear and nonlinear refraction indices of the material, respectively, and $V_{\rm eff}$ denotes the effective volume of the pumped mode. $\eta = \kappa_{\rm ext}/\kappa$ is the coupling efficiency related to coupling (or extrinsic) linewidth $\kappa_{\rm ext}$ and total linewidth κ , where $\kappa = \kappa_{\rm in} + \kappa_{\rm ext}$ and $\kappa_{\rm in}$ is the intrinsic linewidth.

We then consider the dimensionless longitudinal coordinate $\theta = \phi G$ in the anomalous GVD regime ($\beta < 0$), where $G = \sqrt{\frac{1}{2d_2}}$ and $d_2 = D_2/\kappa$ is the dimensionless dispersion. Equation (1) can be further simplified to^[21,22]

$$j\frac{\partial\Psi}{\partial\tau} + \frac{1}{2}\frac{\partial^{2}\Psi}{\partial\theta^{2}} + |\Psi|^{2}\Psi = (-j + \zeta_{0})\Psi + jf, \qquad (3)$$

which is known as the nonlinear Schrödinger equation with damping, detuning, and driving. When the interaction between dispersion and nonlinearity effect is counteracted and the energy in the cavity is balanced, dissipative solitons with invariant envelope along the propagation will be formed in the time domain, and in turn stable OFCs will be generated in the linear frequency domain. In other words, the soliton solution can be the stable solution to Eq. (3), which is similar to the standard normalized NLSE. Based on the similarity between them, in this paper, we use NFT for NLSE to analyze the intracavity signal governed by Eq. (3).

2.2. Periodic nonlinear Fourier transform

There are now a lot of detailed NFT algorithms for NLSE, both for periodic boundary condition^[23] and non-periodic boundary condition, and the latter is usually called vanishing boundary condition.^[24] The basic idea for Eq. (1) is to treat the light propagating in the circular direction inside the resonator as the signal propagating along an unfolded track with periodic boundary conditions,^[25] that is,

$$\Psi(\tau,\phi) \equiv \Psi(\tau,\phi+l), \quad l = 2\pi.$$
(4)

The above periodicity also applies to Eq. (3) by changing the value of *l*. Thus, the NFT algorithm with periodic boundary condition is adopted in this work. According to Ref. [23], where the fast nonlinear Fourier transform (FNFT) for the periodic boundary condition was introduced, the nonlinear frequency domain after FNFT includes two parts: the main spectrum and the auxiliary spectrum.^[26] The auxiliary spectrum is also known as the hyperelliptic mode, which is not within the scope of this paper. Our work focuses only on studies of the main spectrum which consists of discrete eigenvalues that remain invariant during pulse evolution according to the standard NLSE (all terms on the right-hand side of Eq. (3) vanish). The eigenvalue, also known as the nonlinear frequency, is the nonlinear analogue of the frequency given in FT. The difference between the two is that the latter refers to real numbers, while the former is distributed in the complex plane and has real axial symmetry.

For the standard NLSE, its NFT is indeed the IST of Zakharov–Shabat problem (ZSP) which can be expressed in the following form:^[1]

$$\begin{cases} \frac{\partial v_1}{\partial \phi} = -j\lambda v_1 + \Psi v_2, \\ \frac{\partial v_2}{\partial \phi} = -\Psi^* v_1 + j\lambda v_2, \end{cases}$$
(5)

where Ψ is the complex envelope of the time-domain signal, and Ψ^* denotes its conjugate signal. The nonlinear frequency parameter λ is a complex eigenvalue, i.e., $\lambda = \text{Re}(\lambda) + j\text{Im}(\lambda)$, whose real part reflects the frequency and imaginary part reflects the amplitude of the corresponding time-domain signal. More detailed theory can be found in the next section. $v = [v_1 \ v_2]^T$ represents the eigenvector which subjects to the eigenvalue function $Lv = \lambda v$, and

$$L = j \begin{bmatrix} \frac{\partial}{\partial \phi} & -\Psi(\tau, \phi) \\ -\Psi^*(\tau, \phi) & \frac{\partial}{\partial \phi} \end{bmatrix}$$
(6)

is an operator whose eigenvalues are independent of time ϕ . For the fact that we just consider the signal in one period, the solutions of Eq. (5) are denoted as $\Phi(\phi)$ and $\tilde{\Phi}(\phi)$ with the canonical initial conditions (ϕ_0 corresponds to the left boundary of the signal in a period)

$$\Phi(\phi_0) = \begin{bmatrix} 1\\ 0 \end{bmatrix}, \qquad \tilde{\Phi}(\phi_0) = \begin{bmatrix} 0\\ 1 \end{bmatrix}. \tag{7}$$

Define the so-called monodromy matrix

$$M(\lambda) := \left[\Phi(\phi_l) \; \tilde{\Phi}(\phi_l) \right], \tag{8}$$

and its Floquet discriminant $\Delta(\lambda) := \frac{1}{2} \operatorname{tr} M(\lambda)$, where ϕ_l corresponds to the right boundary of the signal in a period according to Eq. (5). Then we can obtain the eigenvalues space \mathcal{M} by the following method when D euqidistant samples of the signal Ψ are given:

$$\mathcal{M} = \left\{ \lambda_k \in \mathbb{C} \left| \Delta(\lambda_k) \in \{\pm 1\}, \frac{\mathrm{d}\Delta}{\mathrm{d}\lambda} \right|_{\lambda = \lambda_k} \neq 0 \right\}.$$
(9)

The eigenvalues space \mathcal{M} consists of discrete complex numbers and can be called as the eigenvalue spectrum in the nonlinear frequency domain. In this work, we only give a brief description of the basic concept of FNFT, and interested readers may refer to Ref. [23] for the detailed derivation process.

Since any signal can be regarded as the combination of soliton component and radiation component, although either of these two can be absent for some specific case,^[26] the eigenvalue spectrum obtained from NFT actually consists of two parts, the eigenvalues corresponding to the soliton component and the eigenvalues corresponding to the radiation component of the signal. Figure 1 shows the eigenvalue spectra of 4 different signals via NFT.



Fig. 1. The eigenvalue spectrum distribution of (a) $\Psi(t) = \operatorname{sech}(t) e^{j0.5}$; (b) $\Psi(t) = \operatorname{sech}(2t) e^{j0.5}$; (c) $\Psi(t) = \operatorname{sech}(t) e^{j\operatorname{sech}(t)}$; (d) $\Psi(t) = 2\operatorname{sech}(2t) e^{j0.5}$. Each inset shows the intensity and phase features of the corresponding time domain signals.

According to Matsuda et al.,^[27] signals in the form of $u(t) = A \operatorname{sech}(At)$ (A is a real number) can be a soliton solution (i.e., stationary solution) of NLSE and have definite eigenvalues $\pm jA/2$. When the phase of such a signal is constant, the eigenvalue occurs at the same position, just as Figs. 1(a)and 1(d) show. There are obvious isolated discrete eigenvalues at ± 0.5 j for signal $\Psi(t) = \operatorname{sech}(t) e^{j0.5}$ and ± 1 j for signal $\Psi(t) = 2 \operatorname{sech}(2t) e^{j0.5}$, and the remaining eigenvalues are all 0. In this sense, we relate such isolated eigenvalues in the eigenvalue spectrum to the soliton component in any signal and the remaining to the radiation component. Considering the physical meaning of eigenvalue λ , a pulse in the form of $u(t) = A \operatorname{sech}(At)$ can also be understood as a signal containing a single nonlinear frequency. Figure 1(b) gives the eigenvalue spectrum of signal $\Psi(t) = \operatorname{sech}(2t) e^{j0.5}$ and all the eigenvalues as shown in the eye-like structure are related to non-soliton signal component because no obvious isolated discrete eigenvalues can be observed. Here we again recall the physical meaning of eigenvalues and it is worth to mention that when

the pulse $u(t) = \operatorname{sech}(At)$ gets narrower by increasing *A*, the eye-like structure expands horizontally. That is to say, a narrower time domain signal corresponds to an increase in the nonlinear frequency component, which is a property similar to that of the linear spectrum. By taking the time-variant phase into consideration, an eye-like structure will also appear near the real axis of λ . As a result, the soliton eigenvalues become below the ideal eigenvalues of $\pm 0.5j$ in Fig. 1(c) compared with Fig. 1(a), which is subjected to the Parseval identity.^[1] In short, different time domain signals show distinct nonlinear frequency domain characteristics via NFT, which is the basis of our work.

3. Eigenvalue spectrum analysis based on periodic NFT

3.1. Case 1

Based on the above property, we can study different signals in the Kerr microresonator, and we use two different sets of parameters for the Kerr OFCs simulation. We first start with a simple case of forming an optical frequency comb with one-way scan of the pump laser. Figure 2 shows the temporal evolution, and the related simulation parameters are $P_{\rm in} =$ 0.755 W, $\lambda_0 = 1550$ nm, $\kappa/2\pi = 287$ MHz, $n_0 = 2.122$, $n_2 =$ 2.4×10^{-19} m²/W, $D_1/2\pi = 44.48$ GHz, $D_2/2\pi = 12.5$ MHz, $V_{\rm eff} = 3.6594 \times 10^{-15}$ m³, and $\eta = 0.9989$.

As can be seen from Fig. 2, different states can be obtained from different detuning via tuning by scanning the pump laser frequency. Clearly, there are four distinct timedomain signals corresponding to the pump laser, Turing pattern, chaos, and soliton state, respectively. It is important to point out that the so-called soliton state here corresponds to the stable solution to Eq. (5) and is different with the soliton component defined early. The periodic FNFT is then performed to these signals and we try to find their corresponding nonlinear spectrum characteristics. Due to the simplicity of the linear frequency domain when there is only pump light in the cavity, we do not consider its nonlinear domain in this paper. Figure 3 shows the time-domain signals (left column) of interest in Fig. 2 and their eigenvalue spectra (right column).



Fig. 2. The temporal evolution diagram inside the Kerr microresonator during one-way scan of the pump laser.

Figure 3(a) describes the time-domain signal of the Turing pattern, which is a state of modulation instability. As shown in Fig. 3(b), its eigenvalue spectrum is strictly symmetric, with respect to both the real axis and the imaginary axis. The eigenvalue spectrum includes several isolated discrete eigenvalue pairs, showing several nonlinear frequency components. According to the specific distribution of the eigenvalues, the Turing pattern temporal signal includes some soliton components defined in our paper because there are eigenvalues on the imaginary axis of the eigenvalue spectrum. The time-domain signal for chaos which can arise in any nonlinear system is shown in Fig. 3(c), and its eigenvalue spectrum (Fig. 3(d)) is also disorderly. When the energy in the cavity reaches an equilibrium, and the dispersion and nonlinear effects cancel each other out, the dissipative soliton forms as shown in Fig. 3(e). Its eigenvalue spectrum includes obvious quasi-continuous eigenvalues corresponding to the radiation component and a pair of soliton discrete eigenvalues, where the imaginary part of the discrete eigenvalues can reflect the amplitude feature of the soliton. It needs to be noted that, sometimes, breathing solitons will occur in the tuning process, that is, the intensity of a temporal soliton will change periodically with time. Under this circumstance, the position variation of the discrete eigenvalues can be utilized to judge the existence of a breathing soliton. Thus, according to different distributions of eigenvalues of these signals, we can easily distinguish different stages during the forming of OFCs in the nonlinear frequency domain.



Fig. 3. The intensity and phase portrait of the intracavity signals (left column) and the corresponding eigenvalue spectra obtained by FNFT (right column). The corresponding normalized detunings are (a) $\zeta_0 = 3.86$, (c) $\zeta_0 = 5.63$, (e) $\zeta_0 = 12.68$.

3.2. Case 2

Next, we consider the more complex case of Kerr OFCs generation with multiple temporal solitons existing in a roundtrip time. The necessary simulation parameters are listed as in Ref. [22] $P_{\rm in} = 2$ W, $\lambda_0 = 1553$ nm, $\kappa/2\pi = 300$ MHz, $n_0 =$ 2.4, $n_2 = 2.4 \times 10^{-19} \text{ m}^2/\text{W}$, $D_1/2\pi = 100 \text{ GHz}$, $D_2/2\pi =$ 2.5 MHz, and $V_{\text{eff}} = 1.0 \times 10^{-15} \text{ m}^3$. Here, we apply the different parameter $\eta = 0.50$ which is the critical coupling efficiency. The resulted temporal evolution diagram is shown in Fig. 4. In this simulation for the generation of Kerr OFCs, the pumped laser frequency is scanned forward and backward, that is, we gradually decrease ω_p and then increase it, which can be observed from the variation of the normalized detuning in the figure. Thus, by combining the tuning process with thermal effect, abundant soliton dynamic processes can be formed in the Kerr microresonator. Similarly, the time-domain signals of interest are extracted from the tuning process, including the Turing pattern, chaos, single soliton, and multi-solitons state.



Fig. 4. The temporal evolution diagram inside the Kerr microresonator by thermal effect and forward and backward tuning of the pump laser.



Fig. 5. The temporal intensity and phase profiles, linear spectrum, and nonlinear eigenvalue spectrum of Turing pattern (left column) and single soliton state (right column). $\zeta_0 = 13.54$ for the Turing pattern generated during the decrease of ω_p , while $\zeta_0 = 23.89$ for the single soliton state generated during the increase of ω_p .

First, we analyze the time-domain signals of Turing pattern and single soliton state. Figure 5 shows the temporal intensity and phase profiles, linear spectrum, and nonlinear eigenvalue spectrum of these two signals. In this case, the generated Turing pattern and soliton signal within an intracavity round-trip time are different from those in case 1, and the obtained eigenvalues are quite different. However, the distribution of eigenvalues follows the same rule, i.e., strictly symmetric discrete eigenvalue pairs for the Turing pattern (see Fig. 5(e)), and combination of a pair of discrete eigenvalues and quasi-continuous spectrum with obvious eye-like structure for the single soliton pulse (see Fig. 5(f)). Figures 5(c)and 5(d) show the linear spectrum. A few frequency components exist in the linear domain for the signal with Turing pattern, which is also known as the modes of the primary comb. Once a stable dissipative soliton pulse is formed, the optical

frequency comb with smooth and stable envelope and wide band will be generated as Fig. 5(d) shows, which is also what we hope to get. Here, we discover that there is a certain correlation between the nonlinear eigenvalue spectrum and the linear spectrum for the fact that the real part of the eigenvalue represents the frequency of the signal. That is, for the signal with Turing pattern, there are only a few frequency components either in the linear or nonlinear frequency domain. For a single soliton pulse, except for a pair of isolated eigenvalues corresponding to the soliton component of the initial signal, all the other nonlinear frequency points are smoothly and uniformly distributed in the nonlinear frequency domain, which is also a property of linear frequency lines distribution.



Fig. 6. The temporal intensity and phase profiles, linear spectrum, and nonlinear eigenvalue spectrum of chaos (left column) and 4-solitons state (right column). $\zeta_0 = 35.16$ for the chaos generated during the decrease of ω_p , while $\zeta_0 = 27.34$ for the 4-solitons state generated during the increase of ω_p .

Then we analyze the chaotic and multi-solitons temporal signals. The temporal characteristics for the chaos and 4solitons state are captured as Fig. 6 shows, as well as their frequency characteristics. The time-domain signal of 4-solitons here is non-periodic in a round trip time, that is, the distance between the 4 solitons is not equal. So, its linear spectrum (Fig. 6(d)) is very similar to that of the chaos (Fig. 6(c)), they both look very noisy, and thus the multi-solitons state may sometimes be mistaken as chaotic states just by identifying the linear spectrum. In essence, however, they are significantly different from each other. That is, the modes of chaotic OFCs are incoherent, so the comb is unstable, whereas the multisolitons comb is generally stable. Based on the NFT analysis of the two time-domain signals in Figs. 6(a) and 6(b), the nonlinear eigenvalue spectra are obtained as shown in Figs. 6(e) and 6(f). As mentioned above, the eigenvalue spectrum of chaos is also chaotic, while the eigenvalue spectrum of the 4-solitons signal is actually ordered because of the strict symmetry of the discrete eigenvalues arisen from its radiation components, which ensures a good distinction between the multisolitons signal and chaotic signal. We can also observe 4 pairs of obvious isolated eigenvalues in Fig. 6(f) corresponding to the signal's soliton components, and the number of solitons in the temporal pulse is also 4. It should be noted that we have carried the proper adjustments to the signal amplitude in the multi-solitons case to capture the signal in the form of soliton components $A \operatorname{sech}(At)$ as close as possible for further NFT analysis.

We also perform the periodic nonlinear Fourier transform analysis of 7-solitons and 9-solitons signals, and obtain the nonlinear eigenvalue spectra as shown in Fig. 7. By adjusting the amplitude of the signal extracted from the temporal evolution, the corresponding soliton eigenvalues and non-soliton eigenvalues separate well in the eigenvalue spectrum. For a more intuitive view, partial eigenvalue spectra for 7-solitons and 9-solitons are presented and thus we just focus on the distribution of isolated soliton eigenvalues. We also find that the number of isolated eigenvalue pairs corresponds to the number of solitons in the time domain, that is, there are 7 pairs of distinct isolated eigenvalues for the 7-solitons temporal signal and 9 pairs for the 9-solitons state.



Fig. 7. The temporal intensity and phase profiles, and nonlinear eigenvalue spectrum of 7-solitons (left column) and 9-solitons states (right column). $\zeta_0 = 37.66$ for 7-solitons state and $\zeta_0 = 58.19$ for 9-solitons state are both generated during the increase of ω_p .

4. Conclusion

In summary, based on the fact that the normalized LLE can be regarded as the NLSE including several other effects, we can apply periodic NFT to the analysis of signals during the generation of OFCs in the Kerr microresonator. We finally find that the signals at different states can be distinguished clearly according to different distributions of the eigenvalue spectrum. For the time-domain signal extracted from a chaotic state, its eigenvalue distribution is also disordered. The eigenvalue spectrum of the Turing pattern is characterized by several strictly symmetric isolated eigenvalue pairs. Specially, the eigenvalue spectrum of the single soliton pulse is composed of a pair of conjugate symmetric isolated eigenvalues and the quasi-continuous eigenvalue spectrum with eye-like structure. Take NFT analysis directly for the multi-solitons time-domain signal in the cavity, the eigenvalue spectrum obtained is complex but ordered. By adjusting the signal amplitude, we can approximately capture the waveforms corresponding to the soliton components. In this way, obvious soliton eigenvalue pairs are obtained by NFT and its number is consistent with the number of solitons formed in the microresonator.

In this work, we have discovered that there is a certain correlation between the nonlinear eigenvalue spectrum and the linear spectrum. However, the accurate correspondence needs to be studied in the future. And we hope to derive the mathematical form of amplitude adjustment in the case of multisolitons from the expression of soliton solution in Kerr microresonator in next work for richer soliton dynamics analysis.

References

- [1] Yousefi M I and Kschischang F R 2014 IEEE Trans. Inf. Theory 60 4312
- [2] Agrawal G P 2007 Nonlinear Fiber Optics, 4th edn. (Academic Press)
- [3] Prilepsky J E, Derevyanko S A, Blow K J, Gabitov I and Turitsyn1 S K
- 2014 Phys. Rev. Lett. 113 013901
- [4] Yousefi M I and Kschischang F R 2014 *IEEE Trans. Inf. Theory* 60 4346
- [5] Wang L N, Liu S Y, Li C Y and He G Q 2017 A Combination of Eigenvalue and Spectral Function Modulation in Nonlinear Frequency Di-

vision Multiplexing, OSA Nonlinear Optics Toptical Meeting (NLO), 17–21 July 2017, Waikoloa, Hawaii, USA

- [6] Liu S Y, Wang L N, Li C Y and He G Q 2017 Spectral Function Modulation based on Nonlinear Fourier Transform, OSA Nonlinear Optics Toptical Meeting (NLO), 17–21 July 2017, Waikoloa, Hawaii, USA
- [7] He G Q, Wang L N, Li C Y, Liu S Y and Hu W S 2017 *Sci. Rep.* **7** 6058
- [8] Son T Le, Aref V and Buelow H 2017 Nat. Photon. 11 570
- [9] Son T Le and Buelow H 2017 *IEEE J. Lightwave Technol.* **35** 3692
- [10] Gui T, Lu C, Lau A P T and Wai P K A 2017 *Opt. Express* **25** 20286
- [11] Goossens J W, Yousefi M I, Jaouën Y and Hafermann H 2017 Opt. Express 25 26437
- [12] Son T Le, Aref V and Buelow H 2018 IEEE J. Lightwave Technol. 36 1296
- [13] Son T Le, Schuh K, Buchali F and Buelow H 2018 Optical Fiber Communication Conference paper W1G.6
- [14] Yangzhang X, Aref V, Son T Le, Buelow H and Bayvel P 2018 arXiv:1806.10367v1
- [15] Wang J, Zhao Y L, Huang X and He G Q 2019 ZTE Commun. 3 17
- [16] Ryczkowski P, Närhi M, Billet C, Merolla J M, Genty G and Dudley J M 2018 Nat. Photon. 12 221

- [17] Kippenberg T J, Gaeta A L, Lipson M and Gorodetsky M L 2018 Science 361 567
- [18] Godey C, Balakireva I V, Coillet A and Chembo Y K 2014 Phys. Rev. A 89 063814
- [19] Chembo Y K and Menyuk C R 2013 Phys. Rev. A 87 053852
- [20] Lamont M R E, Okawachi Y and Gaeta A L 2013 Opt. Lett. 38 3478
- [21] Herr T, Brasch V, Jost J D, Wang C Y, Kondratiev N M, Gorodetsky M L and Kippenberg T J 2014 Nat. Photon. 8 145
- [22] Guo H, Karpov M, Lucas E, Kordts A, Pfeiffer M H P, Brasch V, Lihachev G, Lobanov V E, Gorodetsky M L and Kippenberg T J 2017 *Nat. Phys.* 13 94
- [23] Wahls S and Poor H V 2015 IEEE Trans. Inf. Theory 61 6957
- [24] Yousefi M I and Kschischang F R 2014 IEEE Trans. Inf. Theory 60 4329
- [25] Agha I H, Okawachi Y and Gaeta A L 2009 Opt. Express 17 16209
- [26] Turitsyn S K, Prilepsky J E, Son T Le, Wahls S, Frumin L L, Kamalian M and Derevyanko S K 2017 Optica 4 307
- [27] Matsuda Y, Terauchi H and Maruta A 2014 OptoElectronics and Communication Conference 1016



Chinese Physics B

Volume 29 Number 3 March 2020

SPECIAL TOPIC — Topological semimetals

037101 Single crystal growth, structural and transport properties of bad metal RhSb₂ D S Wu, Y T Qian, Z Y Liu, W Wu, Y J Li, S H Na, Y T Shao, P Zheng, G Li, J G Cheng, H M Weng and J L Luo

SPECIAL TOPIC — Advanced calculation & characterization of energy storage materials & devices at multiple scale

038202 Computational screening of doping schemes for $LiTi_2(PO_4)_3$ as cathode coating materials Yu-Qi Wang, Xiao-Rui Sun, Rui-Juan Xiao and Li-Quan Chen

REVIEW

- 037801 Graphene's photonic and optoelectronic properties A review A J Wirth-Lima, P P Alves-Sousa and W Bezerra-Fraga
- 038701 Astrocyte and ions metabolism during epileptogenesis: A review for modeling studies Meng-Meng Du, Jia-Jia Li, Zhi-Xuan Yuan, Yong-Chen Fan and Ying Wu

RAPID COMMUNICATION

037305 Tailoring electronic properties of two-dimensional antimonene with isoelectronic counterparts

Ye Zhang, Huai-Hong Guo, Bao-Juan Dong, Zhen Zhu, Teng Yang, Ji-Zhang Wang and Zhi-Dong Zhang

GENERAL

- 030201 Lump and interaction solutions to the (3+1)-dimensional Burgers equation Jian Liu and Jian-Wen Wu
- 030202 Lax pair and vector semi-rational nonautonomous rogue waves for a coupled timedependent coefficient fourth-order nonlinear Schrödinger system in an inhomogeneous optical fiber

Zhong Du, Bo Tian, Qi-Xing Qu and Xue-Hui Zhao

030203 Exact solutions of stochastic fractional Korteweg de–Vries equation with conformable derivatives

Hossam A. Ghany, Abd-Allah Hyder and M Zakarya

030301 Construction of Laguerre polynomial's photon-added squeezing vacuum state and its quantum properties

Dao-Ming Lu

030302 Geometric phase of an open double-quantum-dot system detected by a quantum point contact

Qian Du, Kang Lan, Yan-Hui Zhang and Lu-Jing Jiang

(Continued on the Bookbinding Inside Back Cover)

- 030303 Reference-frame-independent quantum key distribution with an untrusted source Jia-Ji Li, Yang Wang, Hong-Wei Li and Wan-Su Bao
- 030304 Quantum speed limit time of a non-Hermitian two-level system Yan-Yi Wang and Mao-Fa Fang
- 030305 Applicability of coupling strength estimation for linear chains of restricted access He Feng, Tian-Min Yan and Yuhai Jiang
- 030306 Optical complex integration-transform for deriving complex fractional squeezing operator Ke Zhang, Cheng-Yu Fan and Hong-Yi Fan
- 030501 Dynamics of the plane and solitary waves in a Noguchi network: Effects of the nonlinear quadratic dispersion S A T Fonkoua, M S Ngounou, G R Deffo, F B Pelap, S B Yamgoue and A Fomethe
- 030502 Dynamics analysis of chaotic maps: From perspective on parameter estimation by metaheuristic algorithm Yue-Xi Peng, Ke-Hui Sun and Shao-Bo He
- 030503 The second Hopf bifurcation in lid-driven square cavity

Tao Wang, Tiegang Liu and Zheng Wang

030504 Dynamical response of a neuron-astrocyte coupling system under electromagnetic induction and external stimulation

Zhi-Xuan Yuan, Pei-Hua Feng, Meng-Meng Du and Ying Wu

- 030505 Hexagonal arrangement of phospholipids in bilayer membranes Xiao-Wei Chen, Ming-Xia Yuan, Han Guo and Zhi Zhu
- 030601 An Yb-fiber frequency comb phase-locked to microwave standard and optical reference Hui-Bo Wang, Hai-Nian Han, Zi-Yue Zhang, Xiao-Dong Shao, Jiang-Feng Zhu and Zhi-Yi Wei

ATOMIC AND MOLECULAR PHYSICS

- 033101 Effects of electron correlation and the Breit interaction on one- and two-electron onephoton transitions in double K hole states of He-like ions ($10 \le Z \le 47$) Xiaobin Ding, Cunqiang Wu, Mingxin Cao, Denghong Zhang, Mingwu Zhang, Yingli Xue, Deyang Yu and Chenzhong Dong
- 033102 Vibronic spectra of aluminium monochloride relevant to circumstellar molecule Jian-Gang Xu, Cong-Ying Zhang and Yun-Guang Zhang
- 033201 Precise measurement of a weak radio frequency electric field using a resonant atomic probe Liping Hao, Yongmei Xue, Jiabei Fan, Jingxu Bai, Yuechun Jiao and Jianming Zhao
- 033401 Enhancement of electron-ion recombination rates at low energy range in the heavy ion storage ring CSRm

Nadir Khan, Zhong-Kui Huang, Wei-Qiang Wen, Shu-Xing Wang, Han-Bing Wang, Wan-Lu Ma, Xiao-Long Zhu, Dong-Mei Zhao, Li-Jun Mao, Jie Li, Xiao-Ming Ma, Mei-Tang Tang, Da-Yu Yin, Wei-Qing Yang, Jian-Cheng Yang, You-Jin Yuan, Lin-Fan Zhu and Xin-Wen Ma ELECTROMAGNETISM, OPTICS, ACOUSTICS, HEAT TRANSFER, CLASSICAL MECHANICS, AND FLUID DYNAMICS

- 034101 Cherenkov terahertz radiation from Dirac semimetals surface plasmon polaritons excited by an electron beam Tao Zhao and Zhenhua Wu
- 034201 Paraxial propagation of cosh-Airy vortex beams in chiral medium Xiao-Jin Yang, Zhen-Sen Wu and Tan Qu
- 034202 Far-field vector-diffraction of off-axis parabolic mirror under oblique incidence Xia-Hui Zeng and Xi-Yao Chen
- 034203 Optical enhanced interferometry with two-mode squeezed twin-Fock states and parity detection

Li-Li Hou, Shuai Wang and Xue-Fen Xu

- 034204 Dynamic manipulation of probe pulse and coherent generation of beating signals based on tunneling-induced inference in triangular quantum dot molecules Nuo Ba, Jin-You Fei, Dong-Fei Li, Xin Zhong, Dan Wang, Lei Wang, Hai-Hua Wang and Qian-Qian Bao
- 034205 A low-noise, high-SNR balanced homodyne detector for the bright squeezed state measurement in 1–100 kHz range

Jin-Rong Wang, Qing-Wei Wang, Long Tian, Jing Su and Yao-Hui Zheng

034206 Evaluation of polarization field in InGaN/GaN multiple quantum well structures by using electroluminescence spectra shift Ping Chen, De-Gang Zhao, De-Sheng Jiang, Jing Yang, Jian-Jun Zhu, Zong-Shun Liu, Wei Liu, Feng

Liang, Shuang-Tao Liu, Yao Xing and Li-Qun Zhang

034207 Eigenvalue spectrum analysis for temporal signals of Kerr optical frequency combs based on nonlinear Fourier transform

Jia Wang, Ai-Guo Sheng, Xin Huang, Rong-Yu Li and Guang-Qiang He

034208 Design of diamond-shape photonic crystal fiber polarization filter based on surface plasma resonance effect

Yongxia Zhang, Jinhui Yuan, Yuwei Qu, Xian Zhou, Binbin Yan, Qiang Wu, Kuiru Wang, Xinzhu Sang, Keping Long and Chongxiu Yu

- 034209 A compact electro-absorption modulator based on graphene photonic crystal fiber Guangwei Fu, Ying Wang, Bilin Wang, Kaili Yang, Xiaoyu Wang, Xinghu Fu, Wa Jin and Weihong Bi
- 034210 The optical nonreciprocal response based on a four-mode optomechanical system Jing Wang
- 034301 High-contrast imaging based on wavefront shaping to improve low signal-to-noise ratio photoacoustic signals using superpixel method Xinjing Lv, Xinyu Xu, Qi Feng, Bin Zhang, Yingchun Ding and Qiang Liu

034302 Second harmonic magnetoacoustic responses of magnetic nanoparticles in magnetoacoustic tomography with magnetic induction

Gepu Guo, Ya Gao, Yuzhi Li, Qingyu Ma, Juan Tu and Dong Zhang

- 034303 Bubble translation driven by pulsation in a double-bubble system Ling-Ling Zhang, Wei-Zhong Chen, Yuan-Yuan Zhang, Yao-Rong Wu, Xun Wang and Guo-Ying Zhao
- 034304 Ultrasonic beam focusing characteristics of shear-vertical waves for contact-type linear phased array in solid Yu-Xiang Dai, Shou-Guo Yan and Bi-Xing Zhang
- 034305 Influence of dynamic tissue properties on temperature elevation and lesions during HIFU scanning therapy: Numerical simulation Xiao Zou, Hu Dong and Sheng-You Qian
- 034501 Simulation-based optimization of inner layout of a theater considering the effect of pedestrians Qing-Fei Gao, Yi-Zhou Tao, Yan-Fang Wei, Cheng Wu and Li-Yun Dong
- 034701 A mass-conserved multiphase lattice Boltzmann method based on high-order difference Zhang-Rong Qin, Yan-Yan Chen, Feng-Ru Ling, Ling-Juan Meng and Chao-Ying Zhang
- 034702 Multi-bubble motion behavior of uniform magnetic field based on phase field model Chang-Sheng Zhu, Zhen Hu and Kai-Ming Wang
- 034703 Electrohydrodynamic behaviors of droplet under a uniform direct current electric field Zi-Long Deng, Mei-Mei Sun and Cheng Yu
- 034704 Interface coupling effects of weakly nonlinear Rayleigh–Taylor instability with double interfaces

Zhiyuan Li, Lifeng Wang, Junfeng Wu and Wenhua Ye

CONDENSED MATTER: STRUCTURAL, MECHANICAL, AND THERMAL PROP-ERTIES

- 036101 Fractional variant of Stokes–Einstein relation in aqueous ionic solutions under external static electric fields Gan Ren and Shikai Tian
- 036102 Synthesis, structure, and properties of Ba₉Co₃Se₁₅ with one-dimensional spin chains Lei Duan, Xian-Cheng Wang, Jun Zhang, Jian-Fa Zhao, Li-Peng Cao, Wen-Min Li, Run-Ze Yu, Zheng Deng and Chang-Qing Jin
- 036201 First-principles investigation on ideal strength of B2 NiAl and NiTi alloys Chun-Yao Zhang, Fu-Yang Tian and Xiao-Dong Ni

CONDENSED MATTER: ELECTRONIC STRUCTURE, ELECTRICAL, MAG-NETIC, AND OPTICAL PROPERTIES

037201 Effect of overdrive voltage on PBTI trapping behavior in GaN MIS-HEMT with LPCVD SiN $_x$ gate dielectric

Tao-Tao Que, Ya-Wen Zhao, Liu-An Li, Liang He, Qiu-Ling Qiu, Zhen-Xing Liu, Jin-Wei Zhang, Jia Chen, Zhi-Sheng Wu and Yang Liu

- 037301 High-mobility SiC MOSFET with low density of interface traps using high pressure microwave plasma oxidation Xin-Yu Liu, Ji-Long Hao, Nan-Nan You, Yun Bai, Yi-Dan Tang, Cheng-Yue Yang and Sheng-Kai Wang
- 037302 Defect engineering on the electronic and transport properties of one-dimensional armchair phosphorene nanoribbons Huakai Xu and Gang Ouyang
- 037303 Influence of $Zr_{50}Cu_{50}$ thin film metallic glass as buffer layer on the structural and optoelectrical properties of AZO films Bao-Qing Zhang, Gao-Peng Liu, Hai-Tao Zong, Li-Ge Fu, Zhi-Fei Wei, Xiao-Wei Yang and Guo-Hua Cao
- 037304 Comparative study on transport properties of N-, P-, and As-doped SiC nanowires: Calculated based on first principles Ya-Lin Li, Pei Gong and Xiao-Yong Fang
- 037306 General principles to high-throughput constructing two-dimensional carbon allotropes Qing Xie, Lei Wang, Jiangxu Li, Ronghan Li and Xing-Qiu Chen
- 037401 Time-dependent Ginzburg-Landau equations for multi-gap superconductors Minsi Li, Jiahong Gu, Long Du, Hongwei Zhong, Lijuan Zhou and Qinghua Chen
- 037501 Three- and two-dimensional calculations for the interface anisotropy dependence of magnetic properties of exchange-spring $Nd_2Fe_{14}B/\alpha$ -Fe multilayers with out-of-plane easy axes

Qian Zhao, Xin-Xin He, Francois-Jacques Morvan, Guo-Ping Zhao and Zhu-Bai Li

037502 Giant low-field magnetocaloric effect in $\text{EuTi}_{1-x}\text{Nb}_xO_3$ (x = 0.05, 0.1, 0.15, and 0.2) compounds

Wen-Hao Jiang, Zhao-Jun Mo, Jia-Wei Luo, Zhe-Xuan Zheng, Qiu-Jie Lu, Guo-Dong Liu, Jun Shen and Lan Li

- 037701 Magnetoelectric effects in multiferroic Y-type hexaferrites $Ba_{0.3}Sr_{1.7}Co_xMg_{2-x}Fe_{12}O_{22}$
- 037702 Surface passivation in n-type silicon and its application in silicon drift detector Yiqing Wu, Ke Tao, Shuai Jiang, Rui Jia and Ve Huand
- 037802 Processes underlying the laser photochromic effect in colloidal plasmonic nanoparticle aggregates

A E Ershov, V S Gerasimov, I L Isaev, A P Gavrilyuk and S V Karpov

INTERDISCIPLINARY PHYSICS AND RELATED AREAS OF SCIENCE AND TECHNOLOGY

038101 A low-dimensional crystal growth model on an isotropic and quasi-free sustained substrate Chenxi Lu, Senjiang Yu, Lingwei Li, Bo Yang, Xiangming Tao and Gaoxiang Ye

- 038102 Low-temperature plasma enhanced atomic layer deposition of large area HfS₂ nanocrystal thin films Ailing Chang, Yichen Mao, Zhiwei Huang, Haiyang Hong, Jianfang Xu, Wei Huang, Songyan Chen and Cheng Li
- 038103 Comparison study of GaN films grown on porous and planar GaN templates Shan Ding, Yue-Wen Li, Xiang-Qian Xiu, Xue-Mei Hua, Zi-Li Xie, Tao Tao, Peng Chen, Bin Liu, Rong Zhang and You-Dou Zheng
- 038104 Coulomb blockade and hopping transport behaviors of donor-induced quantum dots in junctionless transistors Liu-Hong Ma, Wei-Hua Han and Fu-Hua Yang
- 038201 Theoretical study on the relationship between the position of the substituent and the ESIPT fluorescence characteristic of HPIP

Xin Zhang, Jian-Hui Han, You Li, Chao-Fan Sun, Xing Su, Ying Shi and Hang Yin

- 038501 Investigation of gate oxide traps effect on NAND flash memory by TCAD simulation He-Kun Zhang, Xuan Tian, Jun-Peng He, Zhe Song, Qian-Qian Yu, Liang Li, Ming Li, Lian-Cheng Zhao and Li-Ming Gao
- 038502 Enhancement of radiation hardness of InP-based HEMT with double Si-doped plane Ying-Hui Zhong, Bo Yang, Ming-Ming Chang, Peng Ding, Liu-Hong Ma, Meng-Ke Li, Zhi-Yong Duan, Jie Yang, Zhi Jin and Zhi-Chao Wei
- 038503 Numerical and analytical investigations for the SOI LDMOS with alternated high-k dielectric and step doped silicon pillars

Jia-Fei Yao, Yu-Feng Guo, Zhen-Yu Zhang, Ke-Meng Yang, Mao-Lin Zhang and Tian Xia

- 038504 A method to extend wavelength into middle-wavelength infrared based on InAsSb/(Al)GaSb interband transition quantum well infrared photodetector Xuan-Zhang Li, Ling Sun, Jin-Lei Lu, Jie Liu, Chen Yue, Li-Li Xie, Wen-Xin Wang, Hong Chen, Hai-Qiang Jia and Lu Wang
- 038505 Role of remote Coulomb scattering on the hole mobility at cryogenic temperatures in SOI p-MOSFETs

Xian-Le Zhang, Peng-Ying Chang, Gang Du and Xiao-Yan Liu

- 038801 Microstructure evolution and passivation quality of hydrogenated amorphous silicon oxide (a-SiO_x:H) on (100)- and (111)-orientated c-Si wafers Jun-Fan Chen, Sheng-Sheng Zhao, Ling-Ling Yan, Hui-Zhi Ren, Can Han, De-Kun Zhang, Chang-Chun Wei, Guang-Cai Wang, Guo-Fu Hou, Ying Zhao and Xiao-Dan Zhang
- 038901 Simulation study on cooperation behaviors and crowd dynamics in pedestrian evacuation

Ya-Ping Ma and Hui Zhang ERRATUM 039901 Erratum to "Indium doping effect on properties of ZnO nanoparticles synthesized by S Mourad, J El Ghoul, K Omri and K Khirouni